



*Clypeaster campanulatus* forma *campanulatus* (SCHLOTHEIM, 1820): 1a-c: Müllendorf, Bgld (NHMW 2004z0112/0001)

*Clypeaster campanulatus* forma *portentosus* MICHELIN, 1861: 2a-b: Eisenstadt, Bgld (syntype of *C. portentosus* MICHELIN, 1861; NHMW 1846.37.960); 3a-b: Rauchstallbrunngraben, near Baden, NÖ (syntype of *C. alticostatus* MICHELIN, 1861; NHMW 2003z0055/0001)




*Clypeaster campanulatus* forma *partschii* MICHELIN, 1861: 1a-c, 2a-c: Kalksburg, W (syntypes of MICHELIN, 1861; 1: NHMW 1857.38.30; 2: NHMW 1858.III.19)

a: aboral, b: oral, c: left lateral view




*Clypeaster campanulatus* forma *pyramidalis* MICHELIN, 1861: 1a-c: Mitterberg, near Baden, NÖ (holotype of MICHELIN, 1861; NHMW 1843.LIII.1); 2a-c: Rauchstallbrunngraben, near Baden, NÖ (NHMW 1904.VIII.83)

a: aboral, b: oral, c: right lateral view




*Clypeaster campanulatus* forma aff. *gibbosus* DE SERRES, 1829: 1a-c: Kalksburg, W (NHMW 1904.VIII.88)  
*Clypeaster campanulatus* forma aff. *reidii* WRIGHT, 1855: 2a-c: Müllendorf, Bgld (NHMW 1997z0178/1754)

a: aboral, b: oral, c: lateral view (1c: left lateral, 2c: right lateral)




*Clypeaster campanulatus* forma *partschii* MICHELIN, 1861: 1a-b: corroded test revealing plate sutures; Gainfarn, NÖ (NHMW 2003z0003/0020)

*Clypeaster campanulatus* (SCHLOTHEIM, 1820): 2: internal view of aboral surface; Mannersdorf, NÖ (NHMW A3158); 3: horizontal cross section revealing support structures; Kalksburg, W (NHMW A2807); 4: lantern; Mikulov, Czech Republic (NHMW A2812)




*Clypeaster campanulatus* forma aff. *acuminata* DESOR in AGASSIZ & DESOR, 1847: 1a-c: Perchtoldsdorf, W (NHMW 1904.VIII.91)  
*Clypeaster folium* AGASSIZ in AGASSIZ & DESOR, 1847: 2a-c: Stotzing (sandpit Mayer), Bgld (NHMW 2005z0007/0001)

a: aboral, b: oral, c: lateral view (1c: right lateral, 2c: left lateral)




*Clypeaster intermedius* DES MOULINS, 1837: 1a-c: Roggendorf (Schloßthal), NÖ (holotype of *C. subpartschi* SCHAFFER, 1912; KM 32); 2a-c: Grübern, NÖ (HM 644)





*Clypeaster latirostris* MICHELIN, 1861: 1a-b: Gauderndorf (Himmelreich), NÖ (NHMW 1869.III.37); 2a-c: Grübern, NÖ (NHMW 1997z0178/0526); 3: ? Gauderndorf, NÖ (paralectotype, NHMW 1843.II.67); 4: Limberg (Hengl quarry), NÖ (NHMW 2004z0115/0003)





*Clypeaster neudorfensis* LAMBERT, 1928: Devínska Nová Ves, Slovak Republic (AO of *C. melitensis* MICHELIN, 1861; holotype of *C. neudorfensis* LAMBERT, 1928; NHMW 1851.VI.119)





*Clypeaster calabrus* SEGUENZA, 1880: 1a-c, 2: Müllendorf, Bgld (1a-c: GBA 2004/1/4; 2: NHMW 1997z0178/1753a); 3: Kalksburg, W (NHMW 2003z0049/0003)

*Clypeaster barcinensis* ? LAMBERT, 1906: 4: Devínska Nová Ves, Slovak Republic (NHMW 2003z0048/0001)

↩ *Clypeaster calabrus* SEGUENZA, 1880: 1a-b: Kalksburg, W (NHMW 2003z0049/0003); 2a-b, 3a-c: Müllendorf, Bgld (2: NHMW 1997z0178/1753a; 3: NHMW 1997z0178/1753b)